
Situational awareness and forecasting for Norway

FHI COVID-19 modelling team

Week 10, 17 March 2021

Highlights:

- **National epidemiological situation:** Our models indicate that in the present situation, the prevalence of cases and hospitalisations are increasing due to the exponential phase of the epidemic, governed by a high effective reproduction number. This effective reproduction number is stable at 1.4, indicating a high transmission level. In our changepoint model, the effective reproduction number R acting from 22 February until today is estimated to be 1.41 (median, 95% CI 1.24-1.59). This is the highest estimated reproduction number since March 2020. The estimated probability that R is larger than 1 is 100%. Looking back in time since Christmas, the effective reproduction number was estimated to be below 0.7 until January 21; it then started to increase to below 1 until February 7. Then the increase was twice as rapid, to 1.36 (1.12-1.63) until February 21, but has then remained stable at this level until now, essentially. The SMC model estimates the 7-days averaged effective reproduction number during week 9, to be 1.34 (mean, 95% CI 0.99-1.76). In the SMC model, the estimated probability that the daily reproduction number one week ago was above 1 is 97%. The SMC model indicates a stable trend of the averaged effective reproduction number during week 9, which is the last one we have complete data about. The changepoint model and the SMC agree in estimating the same level of the effective reproduction number and that there is no evidence of change at the national level. Due to the delay between transmission and testing/hospitalisation, it is still too early to see the effect of the interventions implemented on March 2 in the report this week (and with more reason the ones implemented in Oslo and Viken at the beginning of week 11). It will be important to follow the development in the next coming days.

Since the start of the epidemic, we estimate that in total, 153.000 (95% CI 140.000- 172.000) persons in Norway have been infected. The current estimate of the detection probability is stably around ~62%. This means that roughly 73.000 cases went undetected in Norway since the start of the epidemic.

- **National forecasting:** During the next three weeks, we expect the situation to worsen in terms of daily incidence of positive cases and new hospitalisations. In one week, on March 21, we estimate ca 2400 new cases per day (median; 95% CI 1500-3500), and a prevalence (total number of presently infected individuals in Norway) of 13.000 (median; 95% CI 8.500-18.000). Compared to our estimates for March 21 made one week ago, these numbers are higher. The number of COVID-19 patients in hospital (daily prevalence) on March 21 is estimated to be 231 (median 95% CI 162-304), and the number of patients on ventilator treatment is estimated to be 37 (median 95% CI 24-53); the corresponding predictions in three weeks (April 4) are 423 (95% CI 267 - 609) and 68 (95% CI 41 - 101). The probability that the surge capacity will exceed 500 ventilator beds is now estimated to be 99%, which is almost certain. There is a probability of 89% that this number will be above 1000. These estimates are significantly worse than the ones we gave a week ago. We have updated our estimates of the lengths of stay in hospital and in the ICU, using data from January 1. These lengths are longer than in the last four months of 2020.
- **Regional epidemiological situation and forecasting:** We estimate the reproduction number since 5 February for all counties, except for Rogaland (1 February), Agder (12 February), Oslo (22

February), Vestfold og Telemark (22 February) and Viken (22 February). In Oslo, the estimate is 1.31 (mean, 95% CI 1.0-1.63). Between February 5 and February 21, the effective reproduction number was significantly higher, 1.8 (1.5-2.1). In Viken, the estimated effective reproduction number since 22 February is 1.3 (mean, 95% CI 1.05 - 1.5), which is similar to the reproduction number in Viken in the period before (5-21 February). We have to wait until next week to possibly see the favourable effect of the interventions of 2 March, and 2-3 additional weeks to see the effect of the interventions from March 14. Vestfold and Telemark has a particularly high estimated effective reproduction number, 1.68 (1.18-2.19). Agder, Rogaland, Nordland and Troms og Finnmark also have effective reproduction numbers above 1. For Innlandet, Møre og Romsdal, Trøndelag and Vestland, estimated effective reproduction numbers appear to be below 1.

Oslo: The number of new cases per day is estimated to be 320 (mean, 95% CI 175 -550) on 21 March and on April 4 to be 500 (95% CI 220-1100). Hospital prevalence in one week is estimated to be 34 (median; 95% CI 16-63), and in three weeks 40 (median; 95% CI 18-78). The situation is worsening.

- **Telenor mobility data, local mobility and foreign visitors:** Inter-municipality mobility, measured as outgoing mobility of mobile phones from each municipality, is stable, though at low levels in all counties. There has been a decrease in mobility in the last week in Innlandet, but otherwise no change from previous weeks. The number of foreign visitors to Norway has stabilised to a high level, with a slight decrease in the last weeks.

What this report contains:

This report presents results based on a mathematical infectious disease model describing the geographical spread of COVID-19 in Norway. The model consists of three layers:

- Population structure in each municipality.
- Mobility data for inter-municipality movements (Telenor mobile phone data).
- Infection transmission model (SEIR-model)

The model produces estimates of the current epidemiological situation at the municipality, county (fylke), and national levels, a forecast of the situation for the next three weeks, and a long term prediction. We run three different models built on the same structure indicated above: (1) a national changepoint model, (2) a regional changepoint model and (3) a national Sequential Monte Carlo model, named SMC model.

How we calibrate the model: The national changepoint model is fitted to Norwegian COVID-19 hospital incidence data from March 10 until yesterday, and data on the laboratory-confirmed cases from May 1 until yesterday. We do not use data before May 1, as the testing capacity and testing criteria were significantly different in the early period.

Note that the results of the national changepoint model are not a simple average or aggregation of the results of the regional changepoint model because they use different data. The estimates and predictions of the regional model are more uncertain than those of the national model. The regional model has more parameters to be estimated and less data in each county; lack of data limits the number of changepoints we can introduce in that model. In the regional changepoint model, each county has its own changepoints and therefore a varying number of reproduction numbers. Counties where the data indicate more variability, have more changepoints.

The national SMC model is also calibrated both to the hospitalisation incidence data (same data as described above) and the laboratory-confirmed cases.

Telenor mobility data: The mobility data account for the changes in the movement patterns between municipalities that have occurred since the start of the epidemic.

How you should interpret the results: The model is stochastic. To predict the probability of various outcomes, we run the model many times in order to represent the inherent randomness.

We present the results in terms of mean values, 95% confidence intervals, medians, and interquartile ranges. We emphasise that the confidence bands might be broader than what we display, because there are several sources of additional uncertainty which we currently do not fully explore: firstly, there are uncertainties related to the natural history of SARS-CoV-2, including the importance of asymptomatic and presymptomatic infection. Secondly, there are uncertainties related to the timing of hospitalisation relative to symptom onset, the severity of the COVID-19 infections by age, and the duration of hospitalisation and ventilator treatment in ICU. We continue to update the model assumptions and parameters in accordance with new evidence and local data as they become available. A full list of all updates can be found at the end of this report.

Estimates of all reproductive numbers are uncertain, and we use their distribution to assure appropriate uncertainty of our predictions. Uncertainties related to the model parameters imply that the reported effective reproductive numbers should be interpreted with caution.

When we forecast beyond today, we use the most recent reproduction number for the whole future, if not explicitly stated otherwise.

In this report, the term patient in ventilator treatment includes only those patients that require either invasive mechanical ventilation or ECMO (Extracorporeal membrane oxygenation).

1 Estimated national reproduction numbers

Calibration of our national changepoint model to hospitalisation incidence data and test data leads to the following estimates provided in table 1. Figure 1 shows the estimated daily number of COVID-19 patients admitted to hospital (1a) and the estimated daily number of laboratory-confirmed SARS-CoV-2 cases (1b), with blue medians and interquartile bands, which are compared to the actual true data, provided in red. The uncertainty captures the uncertainty in the calibrated parameters in addition to the stochastic elements of our model and the variability of other model parameters.

Table 1: Calibration results

Reff	Period
3.09/3.1(2.38-3.89)	From Feb 17 to Mar 14
0.5/0.5(0.4-0.58)	From Mar 15 to Apr 19
0.62/0.61(0.17-0.97)	From Apr 20 to May 10
0.72/0.7(0.16-1.15)	From May 11 to Jun 30
0.88/0.89(0.19-1.6)	From Jul 01 to Jul 31
1.04/1.04(0.74-1.34)	From Aug 01 to Aug 31
0.92/0.92(0.74-1.1)	From Sep 01 to Sep 30
1.28/1.28(1.11-1.46)	From Oct 01 to Oct 25
1.28/1.28(1.1-1.48)	From Oct 26 to Nov 04
0.79/0.79(0.73-0.85)	From Nov 05 to Nov 30
1.08/1.08(1.03-1.13)	From Dec 01 to Jan 03
0.6/0.6(0.52-0.7)	From Jan 04 to Jan 21
0.85/0.85(0.69-1.01)	From Jan 22 to Feb 07
1.36/1.36(1.12-1.63)	From Feb 08 to Feb 21
1.41/1.4(1.24-1.59)	From Feb 22

Median/Mean (95% credible intervals)




Figure 1: A comparison of true data (red) and predicted values (blue) for hospital admissions and test data. The last four data points (black) are assumed to be affected by reporting delay. B) Comparison of our simulated number of positive cases, with blue median and interquartile bands to the actual true number of positive cases, provided in red. The uncertainty captures the uncertainty in the calibrated parameters, in addition to the stochastic elements of our model and the variability of other model parameters. Note that we do not capture all the uncertainty in the test data—our blue bands are quite narrow. This is likely because we calibrate our model parameters on a 7-days moving average window of test data, instead of daily. This is done to avoid overfitting to random daily variation. Moving averages over 7 days are less variable than the daily data.

1.1 National SMC-model: Estimated daily reproduction numbers

In figure 2, we show how our national model fits the national hospital prevalence data (2a) and the daily number of patients receiving ventilator treatment (2b). Those data sources are not used to estimate the parameters, and can therefore be seen as a validation of the model assumptions.




Figure 2: A comparison of true data (red) and predicted values (blue) for hospital and respirator prevalence.

1.1 National SMC-model: Estimated daily reproduction numbers

In the SMC-model, we allow for estimation of a different reproduction number for each day t . To reduce spurious fluctuation, we report a 7-days moving average, $R(t)$, representing the average reproduction number for the whole week before day t . However, until March 8 we keep the reproduction number constant. By assuming a time varying reproduction number $R(t)$, we can detect changes without introducing explicit changepoints. Thus, we can easier detect unexpected changes.

The SMC model uses the daily number of new admissions to hospital and the daily number of positive and negative lab-confirmed tests, to estimate all its parameters. Because of the time between infection and the possibility to be detected as positive by a test, and because of a delay in reporting tests, the data contain information on the transmissibility until a week before the end of the data (today).

The parameters π_0 and π_1 related to the probability to detect a positive case by testing are estimated off-line.

The figure below shows the SMC estimate of the 7-day-average daily reproduction number $R(t)$ from the start of the epidemic in Norway and until today. In the figure we plot the 95% confidence interval and quantiles of the estimated posterior distribution of $R(t)$.

1.1 National SMC-model: Estimated daily reproduction numbers




Figure 3: $R(t)$ estimates using a Sequential Monte Carlo approach calibrated to hospitalisation incidence and test data. The large uncertainty during the last 7 days reflects the lack of available data due to the transmission delay, test delay, time between symptoms onset and hospitalisation. The green band shows the 95% posterior credibility interval. As we use test data only from 1 August, the credibility interval becomes more narrow thereafter.

2 National estimate of cumulative (total) number of infections


The national changepoint model estimates the total number of infections and the symptomatic cases that have occurred (Table 2).

Figure 4a shows the modelled expected daily incidence (blue) and the observed daily number of laboratory-confirmed cases (red). When simulating the laboratory-confirmed cases, we also model the detection probability for the infections (both symptomatic, presymptomatic and asymptomatic), Figure 4b. There are two differences between this estimate of the detection probability and the previous one provided in figure 4a. In figure 4b, we calibrate our model to the true number of positive cases, instead of using the test data directly. Furthermore, in figure 4a we use a parametric model to estimate the detection probability that depends on the true total number of tests performed.

Table 2: Estimated cumulative number of infections, 2021-03-14

Region	Total	No. confirmed	Fraction reported	Min. fraction
Norway	153098 (133916; 172837)	80440	53%	47%

Fraction reported=Number confirmed/number predicted; Minimal fraction reported=number confirmed/upper CI



(a) Number of laboratory-confirmed cases vs model-based estimated number of new infected individuals
(b) Estimated detection probability for an infected case per calendar day

Figure 4

3 National 3-week predictions: Prevalence, Incidence, Hospital beds and Ventilator beds

The national changepoint model estimates the prevalence and daily incidence of infected individuals (asymptomatic, presymptomatic and symptomatic) for the next three weeks, aggregated to the whole of Norway (table 3). In addition, the table shows projected national prevalence of hospitalised patients (hospital beds) and prevalence of patients receiving ventilator treatment (ventilator beds). The projected epidemic and healthcare burden are illustrated in figure 5.

Table 3: Estimated national prevalence, incidence, hospital beds and ventilator beds. Median/Mean (CI)

	1 week prediction (Mar 21)	2 week prediction (Mar 28)	3 week prediction (Apr 04)
Prevalence	12852/12598 (8479-17781)	17532/17130 (10445-26186)	23861/23042 (12956-38299)
Daily incidence	2387/2324 (1505-3511)	3246/3163 (1849-4925)	4420/4243 (2266-7347)
Hospital beds	231/228 (162-304)	313/309 (217-437)	423/412 (267-609)
Ventilator beds	37/36 (24-53)	50/50 (32-72)	68/67 (41-101)




Figure 5: National 3 week predictions for incidence (top left), prevalence (bottom left), hospital beds (top right) and ventilator beds (bottom right)

4 National long-term scenarios: Prevalence, Hospital beds and Ventilator beds

NB: Vaccination of individuals is not included in these predictions.

Results from 12-month scenario of the calibrated national changepoint model, showing expected prevalence (Figure 6a), hospital beds (Figure 6b) and ventilator beds (Figure 6c), in the case where the transmissibility stays the same as today. The figures are made using the 200 candidate models, where the reproductive numbers are varying according to their estimated uncertainty as of today. The confidence intervals shown in the plots are two-tailed around the median, and therefore the upper 95% level shows the 97.5% boundary. Note that age-specific attack rate after 21 days of projection is assumed to follow the demography in each county, instead of being informed by the current age-distribution of the laboratory-confirmed cases.




Figure 6: Long-term predictions for prevalence (a), hospital beds (b) and ventilator beds (c)

The probability of a surge capacity need above **500 ICU** ventilator beds is **99 %**. We estimate the probability of a surge capacity need above **1000 ICU** beds to be equal to **89 %**.

5 National scenario-based long-term predictions: Hospital beds and Ventilator beds

NOTE: In this section, we do not include any vaccinations, as planned in the next weeks and months. The scenarios are therefore not useful anymore. We will soon substitute them with scenarios which instead include planned vaccinations.

Here we show how the epidemic estimated from the national changepoint model will develop under three assumed epidemiological scenarios, by fixing the effective reproduction number to be 1.1, 1.2 or 1.3, from today. We show the daily number of COVID-19 patients in hospital, including patients receiving ventilator treatment, (Figure 7, and the daily number of patients on ventilator treatment, figure 8. Note that age-specific attack rate after 21 days of projection is assumed to follow the demography in each county, instead of being informed by the current age-distribution of the laboratory-confirmed cases. Additional information about the total attack rate (cumulative incidence) and healthcare burden and surge capacity for these scenarios are provided in Table 4.

Table 4: Predicted numbers of total infected, total number of hospitalisations, total number needing ventilator treatment, and the predicted peak number in ward (not in respirator), hospitalised (both with and without ventilator treatment) and ventilated treatments based on three different scenarios with R effective equal to 1.1, 1.2 and 1.3.

	Reff=1.1	Reff=1.2	Reff=1.3
Total:			
Attack rate (infected)	956 000(917 000 - 1 000 000)	1 630 000(1 610 000 - 1 650 000)	2 210 000(2 200 000 - 2 230 000)
Hospitalisations	28 500(27 300 - 29 800)	49 800(49 100 - 50 500)	68 100(67 300 - 68 900)
Patients on ventilator	3 070(2 930 - 3 210)	5 320(5 190 - 5 440)	7 250(7 110 - 7 410)
At peak:			
Hospital beds, excl. vent.	728(638 - 855)	1 890(1 790 - 2 020)	3 670(3 530 - 3 790)
Hospital beds, incl. vent	888(777 - 1 040)	2 330(2 200 - 2 470)	4 500(4 340 - 4 660)
Ventilator beds	177(153 - 204)	450(417 - 485)	861(819 - 907)




Figure 7: Predicted number of COVID-19 patients in hospital based on three different scenarios with R effective equal to 1.1, 1.2 and 1.3. Shaded areas show interquartile range and 95% confidence interval around the median.




Figure 8: Predicted number of COVID-19 patients needing ventilator treatment based on three different scenarios with R effective equal to 1.1, 1.2 and 1.3. Shaded areas show interquartile range and 95% confidence interval around the median.

6 Estimated regional reproduction numbers

Calibration of our regional changepoint model to hospitalisation incidence data and test data leads to the following estimates for current regional reproduction numbers by county (Table 5). A full list of all regional reproduction numbers can be found at the end of the report.

Below we show the estimated daily number of COVID-19 patients admitted to hospital and the estimated daily number of laboratory-confirmed SARS-CoV-2 cases for each county. Model estimates are shown with blue medians and interquantile bands, which are compared to the actual true data, provided in red. The blue bands describe the uncertainty in the calibrated parameters, in addition to the stochastic elements of our model. Last four data points are shown in black as they may be affected by reporting delay.

Table 5: Estimated current regional reproduction numbers

R	Parameter	County	From	Pr(R>1)
1.31 (1.02-1.63)	R9	Oslo	2021-02-22	0.98
1.19 (0.6-1.6)	R7	Rogaland	2021-02-01	0.81
0.52 (0.04-1.16)	R7	Møre og Romsdal	2021-02-05	0.07
1.22 (0.75-1.65)	R9	Nordland	2021-02-05	0.84
1.3 (1.05-1.59)	R9	Viken	2021-02-22	0.99
0.48 (0.08-0.89)	R9	Innlandet	2021-02-05	0
1.68 (1.18-2.19)	R9	Vestfold og Telemark	2021-02-22	1
1.13 (0.73-1.47)	R8	Agder	2021-02-12	0.76
0.74 (0.39-1.06)	R9	Vestland	2021-02-05	0.06
0.72 (0.1-1.3)	R7	Trøndelag	2021-02-05	0.22
1.25 (0.64-1.79)	R7	Troms og Finnmark	2021-02-05	0.82

Mean and 95% credible intervals
















Figure 9: The map shows the direction of the trend in incidence in the counties based on the latest reproduction numbers shown in the other chart. The trend is increasing if the probability that the latest reproduction number is above one is above 95%, the trend is likely increasing if this probability is between 80% and 95%, the trend is uncertain if the probability is between 20% and 80%, the trend is likely decreasing if the probability is between 5% and 20% and is decreasing if the probability that the latest R is above one is less than 5%.






Estimated vs observed hospital incidence data by county:





Estimated and observed lab-confirmed test data by county:





7 Regional 3-week predictions: Cumulative (total) incidence and Prevalence

Below is shown the estimated short-term forecasting of total incidence of infected individuals (table 6), daily incidence (table 7) and prevalence (table 8) for each county.

Table 6: Estimated cumulative number of infections from 2021-01-01 to 2021-03-14

Region	Total	No. confirmed	Fraction reported	Min. fraction
Oslo	13639 (9397; 18989)	8472	62%	45%
Rogaland	1613 (908; 2886)	1317	82%	46%
Møre og Romsdal	377 (178; 811)	330	88%	41%
Nordland	892 (262; 2203)	544	61%	25%
Viken	17442 (12287; 23763)	10066	58%	42%
Innlandet	1545 (937; 2419)	724	45%	30%
Vestfold og Telemark	4593 (2779; 6764)	1944	42%	29%
Agder	1186 (602; 2283)	1185	100%	52%
Vestland	1878 (974; 3271)	1324	71%	40%
Trøndelag	1284 (600; 2439)	821	64%	34%
Troms og Finnmark	369 (168; 803)	363	98%	45%

Fraction reported=Number confirmed/number predicted; Minimal fraction reported=number confirmed/upper CI

Table 7: Predicted incidence per day: Median/Mean (CI)

Region	1 week prediction (21 Mar)	2 weeks prediction (28 Mar)	3 weeks prediction (04 Apr)
Agder	45/49 (17-104)	56/64 (19-148)	73/87 (21-231)
Innlandet	57/61 (26-115)	54/60 (24-129)	62/71 (26-158)
Møre og Romsdal	6/7 (1-19)	3/5 (0-19)	3/5 (0-24)
Nordland	26/32 (6-99)	32/42 (6-140)	38/55 (6-196)
Oslo	304/317 (175-546)	375/403 (191-792)	472/515 (219-1107)
Rogaland	43/50 (13-134)	50/63 (10-193)	59/80 (8-277)
Troms og Finnmark	18/22 (4-65)	23/31 (3-106)	29/45 (4-173)
Trøndelag	16/19 (4-50)	14/19 (2-63)	13/20 (2-78)
Vestfold og Telemark	267/296 (115-609)	418/479 (143-1184)	623/755 (176-2118)
Vestland	39/42 (17-84)	36/41 (13-94)	37/42 (12-101)
Viken	661/703 (389-1204)	812/876 (418-1694)	988/1099 (452-2357)

Table 8: Predicted prevalence. Number of infectious individuals (asymptomatic plus pre-symptomatic plus symptomatic) per day. Median/Mean and 95 perc. CI for three weeks prediction.

Region	21 Mar	28 Mar	04 Apr	low CI, 04 Apr	high CI, 04 Apr
Agder	258/278	332/367	472/532	174	1238
Innlandet	387/406	373/402	493/540	238	1069
Møre og Romsdal	42/47	29/39	26.5/38	7	129
Nordland	147/177	180/229	223/300	52	1041
Oslo	1670/1747	2045.5/2184	2492.5/2711	1213	5646
Rogaland	252/288	305.5/362	369/468	98	1480
Troms og Finnmark	105/125	141/176	190/256	52	883
Trøndelag	115.5/129	102.5/127	117/148	39	426
Vestfold og Telemark	1381.5/1495	2131/2398	3126.5/3704	975	9452
Vestland	259/274	265/291	323/356	152	769
Viken	3792.5/3954	4575/4882	5545/6075	2746	12358

8 Regional 3-week predictions: Hospital beds and ventilator beds

Below is shown the estimated short-term forecasting of expected hospital prevalence (table 9) and patients on ventilator treatment for each county (table 10).

Table 9: Number of hospitalisation beds occupied by Covid-19 patients: Median/Mean (CI)

Region	1 week prediction (21 Mar)	2 weeks prediction (28 Mar)	3 weeks prediction (04 Apr)
Agder	3/4 (0-12)	5/6 (0-15)	7/7 (1-20)
Innlandet	5/5 (1-13)	7/7 (1-18)	7/8 (1-19)
Møre og Romsdal	0/0 (0-3)	0/1 (0-4)	0/1 (0-4)
Nordland	1/2 (0-9)	2/3 (0-10)	2/4 (0-15)
Oslo	34/34 (16-59)	35/36 (16-63)	39/41 (18-78)
Rogaland	2/3 (0-10)	3/4 (0-14)	5/6 (0-20)
Troms og Finnmark	0/1 (0-5)	1/1 (0-7)	1/2 (0-9)
Trøndelag	1/1 (0-7)	2/2 (0-8)	2/3 (0-8)
Vestfold og Telemark	21/22 (8-42)	30/32 (10-65)	43/49 (14-112)
Vestland	2/3 (0-9)	3/4 (0-12)	4/5 (0-13)
Viken	51/53 (28-86)	65/67 (34-115)	80/85 (40-154)

Table 10: Number of ICU beds occupied by Covid-19 patients: Median/Mean (CI)

Region	1 week prediction (21 Mar)	2 weeks prediction (28 Mar)	3 weeks prediction (04 Apr)
Agder	0/1 (0-3)	1/1 (0-3)	1/1 (0-4)
Innlandet	1/1 (0-3)	1/1 (0-4)	1/1 (0-4)
Møre og Romsdal	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Nordland	0/0 (0-2)	0/0 (0-2)	0/1 (0-3)
Oslo	6/6 (2-12)	6/7 (2-13)	7/7 (2-15)
Rogaland	0/0 (0-2)	0/1 (0-3)	1/1 (0-4)
Troms og Finnmark	0/0 (0-1)	0/0 (0-2)	0/0 (0-2)
Trøndelag	0/0 (0-2)	0/0 (0-2)	0/0 (0-2)
Vestfold og Telemark	3/3 (0-8)	5/5 (1-11)	7/7 (1-17)
Vestland	0/1 (0-2)	0/1 (0-3)	1/1 (0-3)
Viken	8/9 (3-17)	11/11 (4-21)	13/14 (5-26)

Table 11: Trend analysis for the last 14 days

County	Average daily increase last 14 days		Doubling Time (days)	
	Hospitalisations	Cases	Hospitalisations	Cases
Agder	Not enough data	3.9 (-2.1, 10.2) %	Not enough data	18.3 (-33.4, 7.2)
Innlandet	Not enough data	4.8 (-1, 11.1) %	Not enough data	14.8 (-68.7, 6.6)
Møre og Romsdal	Not enough data	8.8 (-1.8, 21.2) %	Not enough data	8.2 (-37.2, 3.6)
Nordland	Not enough data	9.7 (0.1, 20.3) %	Not enough data	7.5 (469.5, 3.8)
Norge	7 (2, 12.4) %	4.5 (3.6, 5.5) %	10.2 (35.8, 5.9)	15.6 (19.4, 13)
Oslo	9.6 (1.2, 19.2) %	5.2 (3.6, 6.7) %	7.5 (56, 3.9)	13.7 (19.4, 10.6)
Rogaland	Not enough data	16.7 (7.9, 26.6) %	Not enough data	4.5 (9.1, 2.9)
Troms og Finnmark	Not enough data	48.6 (25.7, 83.7) %	Not enough data	1.8 (3, 1.1)
Trøndelag	Not enough data	10.3 (1.7, 20.2) %	Not enough data	7.1 (40.2, 3.8)
Vestfold og Telemark	Not enough data	-3.7 (-7.2, -0.1) %	Not enough data	-18.3 (-9.3, -534.9)
Vestland	-8.3 (-25.7, 10.6) %	-3 (-7.7, 2) %	-8 (-2.3, 6.9)	-23.1 (-8.6, 34.9)
Viken	6.8 (-1.3, 16) %	4.1 (1.7, 6.4) %	10.5 (-52, 4.7)	17.4 (40, 11.1)

9 14-day trend analysis of confirmed cases and hospitalisations by county

To estimate recent trends in hospitalisation and number of positive tests, we present results in table 11 based on a negative binomial regression where we account for weekend effects. We exclude the last three days to avoid problems of reporting delay and fit the model using data from 17 days to 3 days before the current date. We fit a separate trend model for confirmed cases and for hospital incidence. We only fit a trend model if there has been more than 5 cases or hospitalisations in the 14-day period.

10 Scenario-based short-term predictions for Oslo:

Oslo has experienced increasing infection levels in the last months. Rising case numbers can lead to less efficient contact tracing due to a lack of resources. This, in turn, can cause the reproductive number to increase. To explore the short-term consequences of a less effective contact tracing in Oslo, we compare projections of the regional changepoint model, where the current reproduction number in Oslo is increased to 1.40; 1.45 and 1.50 from today, respectively. In these scenarios we assume no change to the reproductive numbers in the other counties. Table 12 and Figure compares these projected scenarios with a projection of the current epidemiological situation in Oslo.

Table 12: 4 week predictions in Oslo: Prevalence and Incidence (mean/median(CI))

Scenario	Prevalence	Incidence
Current	3234/3074 (1873-5737)	626/594 (355-1100)
R=1.40	3451/3286 (2062-5881)	670/628 (394-1120)
R=1.45	3707/3512 (2173-6332)	727/695 (409-1254)
R=1.50	3983/3788 (2438-6787)	788/736 (486-1347)





Figure 20: Future predictions for Oslo assuming the reproductive number will remain constant vs alternative scenarios. Confidence intervals correspond to "current scenario". Parameters showed are: Hospitalisations (top left), On respirator (top right), Incidence (bottom left) and Prevalence (bottom right).


11 Regional long-term predictions

11.1 Incidence

Predicted incidence (asymptomatic, pre-symptomatic and symptomatic) of the calibrated regional change-point model for each county per day, with confidence intervals.




11.2 Hospitalisations




11.2 Hospitalisations

Estimated prevalence of COVID-19 patients in hospital, including patients receiving ventilator treatment.

11.2 Hospitalisations



11.2 Hospitalisations



12 Mobility data

Number of trips out from each municipality during each day is based on Telenor mobility data. We observed a large reduction in inter-municipality mobility in March 2020 (with minimum reached on Tuesday 17 March 2020), and thereafter we see an increasing trend in the mobility lasting until vacation time in July. The changes in mobility in July coincides with the three-week "fellesferie" in Norway, and during August the mobility resumes approximately the same levels as pre-vacation time. There is however a significant regional variation.

The reference level is set to 100 on March 2nd 2020 for all the figures in this section, and we plot the seven-day, moving average of the daily mobility. Figure 21 shows an overview of the mobility since March 2020 for the largest municipalities in each county, and Figure 22 shows the total mobility out from all municipalities in each county, including Oslo. Figure 23 and 24, zooms in on mobility from December 14 2020, for municipalities and counties, respectively.




Figure 21: Mobility for selected municipalities since March 2020: Nationally (Norge), Stavanger (1103), Ålesund (1507), Bodø (1804), Bærum (3024), Ringsaker (3411), Sandefjord (3804), Kristiansand (4204), Bergen (4601), Trondheim (5001), Tromsø (5401).




Figure 22: Mobility for fylker since March 2020: Oslo (03), Rogaland (11), Møre og Romsdal (15), Nordland (18), Viken (30), Innlandet (34), Vestfold og Telemark (38), Agder (42), Vestland (46), Trøndelag (50), Troms og Finmark (54).




Figure 23: Zoom: Mobility from December 14, 2020 and onwards: Nationally (Norge), Stavanger (1103), Ålesund (1507), Bodø (1804), Bærum (3024), Ringsaker (3411), Sandefjord (3804), Kristiansand (4204), Bergen (4601), Trondheim (5001), Tromsø (5401).




Figure 24: Zoom: Mobility from December 14, 2020 and onwards, per fylker: Oslo (03), Rogaland (11), Møre og Romsdal (15), Nordland (18), Viken (30), Innlandet (34), Vestfold og Telemark (38), Agder (42), Vestland (46), Trøndelag (50), Troms og Finnmark (54).

	7	8	9	10	11
Norge	72.2	73.6	75.7	77.1	73.7
Stavanger	68.7	69.5	70.9	69.8	71.7
Ålesund	84.9	88.0	86.1	84.1	85.2
Bodø	82.6	82.7	80.0	79.0	76.7
Bærum	48.1	50.1	46.8	47.3	47.9
Ringsaker	77.5	79.7	86.4	86.8	81.5
Sandefjord	62.8	63.4	61.7	64.7	57.9
Kristiansand	74.7	76.0	75.6	70.4	69.6
Bergen	67.5	65.9	70.1	74.8	71.3
Trondheim	75.6	75.1	83.2	76.6	79.2
Tromsø	75.5	79.2	97.3	76.0	78.5

Table 13: Municipalities

	7	8	9	10	11
Oslo	45.1	47.3	47.6	45.9	44.9
Rogaland	75.5	76.1	76.2	75.4	77.4
Møre og Romsdal	91.7	93.4	91.6	90.0	90.1
Nordland	89.9	90.3	90.4	89.6	89.8
Viken	62.8	64.9	65.4	67.7	64.8
Innlandet	90.8	93.4	107.5	107.8	96.4
Vestfold og Telemark	72.5	73.8	73.5	78.8	69.2
Agder	85.2	85.2	82.6	88.2	80.6
Vestlandet	77.0	76.3	78.8	83.8	81.4
Trøndelag	84.0	85.9	91.9	89.8	87.4
Troms og Finnmark	85.0	86.5	91.4	86.2	87.3

Table 14: Counties

Weekly mobility for Norway and selected municipalities is displayed in Table 13 and mobility for counties is displayed in Table 14. The percentages in the tables are to be interpreted towards the reference level of 100 for week 10 in March 2020. The color-coding encodes the following: 'Green' monotonic decrease in mobility, 'Yellow' almost monotonic decrease or flat mobility trend, 'Red' increasing mobility.

12.1 Foreign roamers on Telenor's network in Norway

12.1 Foreign roamers on Telenor's network in Norway

An analysis of foreign roamers in Norway from January 2020 has been carried out, to better understand the potential virus importation. In Figure 25 the total number of roamers per day per county are displayed. We can see an approximate 40% drop in the number of visiting roamers after the lock-down in March 2020. The number of visiting roamers recover during the Summer of 2020, and there is a spike of visitors in August followed by a drop again. During October and November 2020 the levels of visiting, foreign roamers to Norway have reached quite high levels, just 10% short of the all-year high for 2020, and Oslo and Viken have seen big increases in visitors. There is a reduction in visitors during Christmas, and in January 2021 we see an increasing trend again.

Figure 26 showcases the levels of roamers from the following countries: Poland, Lithuania, Sweden, Netherlands, Denmark, Latvia, Germany, Spain, Finland and the rest of the world. These levels represent the total number of foreign, visiting roamers from each of the countries per day in Norway, since November 30 2020.




Figure 25: The total number of foreign roamers in Norway broken down on different fylker: Oslo (3), Rogaland (11), Møre og Romsdal (15), Nordland (18), Viken (30), Innlandet (34), Vestfold og Telemark (38), Agder (42), Vestland (46), Trøndelag (50), Troms og Finnmark (54).

12.1 Foreign roamers on Telenor's network in Norway






Figure 26: National overview of total number of foreign, visiting roamers from Poland, Lithuania, Sweden, Netherlands, Denmark, Latvia, Germany, Spain, Finland and the rest of the world.

12.2 Foreign roamers per county (fylke) in Norway

12.2 Foreign roamers per county (fylke) in Norway



12.2 Foreign roamers per county (fylke) in Norway



13 Methods

13.1 Model

We use a metapopulation model to simulate the spread of COVID-19 in Norway in space and time. The model consists of three layers: the population structure in each municipality, information about how people move between different municipalities, and local transmission within each municipality. In this way, the model can simulate the spread of COVID-19 within each municipality, and how the virus is transported around in Norway.

13.1.1 Transmission model

We use an SEIR (Susceptible-Exposed-Infected-Recovered) model without age structure to simulate the local transmission within each area. Mixing between individuals within each area is assumed to be random. Demographic changes due to births, immigration, emigration and deaths are not considered. The model distinguishes between asymptomatic and symptomatic infection, and we consider presymptomatic infectiousness among those who develop symptomatic infection. In total, the model consists of 6 disease states: Susceptible (S), Exposed, infected, but not infectious (E_1), Presymptomatic infected (E_2), Symptomatic infected (I), Asymptomatic infected (I_a), and Recovered, either immune or dead (R). A schematic overview of the model is shown in figure 29.




Figure 29: Schematic overview of the model.

13.2 Movements between municipalities:

We use 6-hourly mobility matrices from Telenor to capture the movements between municipalities. The matrices are scaled according to the overall Telenor market share in Norway, estimated to be 48%. Since week 8, we use the actual daily mobility matrices to simulate the past. In this way, alterations in the mobility pattern will be incorporated in our model predictions. To predict future movements, we use the

13.3 Healthcare utilisation

latest weekday measured by Telenor, regularised to be balanced in total in- and outgoing flow for each municipality.

13.3 Healthcare utilisation

Based on the estimated daily incidence data from the model and the population age structure in each municipality, we calculated the hospitalisation using a weighted average. We correct these probabilities by a factor which represents the over or under representation of each age group among the lab confirmed positive cases. The hospitalisation is assumed to be delayed relative to the symptom onset. We calculate the number of patients admitted to ventilator treatment from the patients in hospital using age-adjusted probabilities and an assumed delay.

13.4 Seeding

At the start of each simulation, we locate 5.367.580 people in the municipalities of Norway according to data from SSB per January 1. 2020. All confirmed Norwegian imported cases with information about residence municipality and test dates are used to seed the model, using the data available until yesterday. For each case, we add an additional random number of infectious individuals, in the same area and on the same day, to account for asymptomatic imported cases who were not tested or otherwise missed. We denote this by the amplification factor.

13.5 Calibration

Estimation of the parameters of the model: the reproduction numbers, the amplification factor for the imported cases, the parameters of the detection probability and the delay between incidence and test, is done using Sequential Monte Carlo Approximate Bayesian Computation (SMC-ABC), as described in Engebretsen et al. (2020): <https://royalsocietypublishing.org/doi/10.1098/rsif.2019.0809>, where the algorithm can be found in the supplement.

The idea behind ABC is to try out different parameter sets, simulate using these, then compare how much the simulations deviate from the observations in terms of summary statistics. We thus test millions of combinations of the different reproductive numbers, the amplification factor, and the parameters for the positive tests, to determine the ones that lead to the best fits to the true number of hospitalised individuals, from March 10 2020 until the last available data point, and the laboratory-confirmed COVID-19 cases from May 1 until the latest available data point.

In the ABC procedure we thus use two summary statistics, one is the distance between the simulated hospitalisation incidence and the observed incidence, and the other is the distance between the observed number of laboratory-confirmed cases and the simulated ones. As the two summary statistics are not on the same scale, we use two separate tolerances in the ABC-procedure, ensuring that we get a good fit to both data sources.

13.5.1 Calibration to hospitalisation data

In order to calibrate to the hospitalisation data, we need to simulate hospital incidence. The details on how we simulate hospitalisations are described in Section 13.3, using the parameters provided in Section 14, which are estimated from individual-level Norwegian data, and updated regularly. As our distance measure, we calculate the squared distance over each time point and each county.

13.5.2 Calibration to test data

We include the laboratory-confirmed cases in the calibration procedure, as these contain additional information about the transmissibility, and the delay between transmission and testing is shorter than the delay between transmission and hospitalisation. Therefore, we simulate also the number of detected positive cases in our model. We assume that the number of detected positive cases can be modelled as a binomial process of the simulated daily total incidence of symptomatic and asymptomatic cases, with a

13.6 Specifications for the national changepoint model

success probability π_t , which changes every day. We also assume a delay d between the day of test and the day of transmission.

The data on the number of positive cases are more difficult to use, as the test criteria and capacity have changed multiple times. We take into account these changes by using the total number of tests performed on each day, as a good proxy of capacity and testing criteria. Moreover, we choose not to calibrate to the test data before May 1, because the test criteria and capacity were so different in the early period. The detection probability is modelled as

$$\pi_t = \exp(\pi_0 + \pi_1 \cdot k_t) / (1 + \exp(\pi_0 + \pi_1 \cdot k_t)),$$

where k_t is the number of tests actually performed on day t , and π_0 and π_1 are two parameters that we estimate, assuming positivity of π_1 . We also estimate the delay d . We choose to use a 7-days backwards moving average for the covariate k_t . To calculate the distance between the observed number of positive tests and the simulated ones we also use a 7-days backwards moving average. We do this to take into account potential day-of-the-week-effects. For example, it could well be that the testing criteria are different on weekends and weekdays. However, using instead the number of tests and calibrating on a daily basis would lead to a larger day-to-day variance. This is likely why we find that the uncertainty in the simulated positive cases seems somewhat too low, and that we do not capture all the variance in the daily test data. Moreover, the binomial assumption could be too simple, and a beta-binomial distribution would allow more variance. A limitation of our current model for the detection probability, is that we only capture the changes in the test criteria that are captured in the total number of tests performed.

13.6 Specifications for the national changepoint model

In the national changepoint model, we assume a first reproduction number R_0 until March 14, a second reproduction number R_1 until April 19, a third reproduction number R_2 until May 10, a fourth reproduction number R_3 until June 30, R_4 until July 31, R_5 until August 31, R_6 from September 1 until September 30, R_7 from October 1 until October 26, R_8 until November 4, R_9 from November 5th until November 30th, R_{10} from December 1st until January 4, a twelfth reproduction number R_{11} from January 4 until January 21, a thirteenth reproduction number from January 22 to February 7 and a fourteenth reproduction number from February 8. This last reproduction number is used for the future. The changepoints follow the changes in restrictions introduced. In the calibration procedure, we obtain 200 parameter sets that we use to represent the distributions of parameters.

After we have obtained the estimated parameters, we run the model with these 200 parameter sets again, from the beginning until today, plus three weeks into the future (or for an additional year). In this way, we obtain different trajectories of the future, allowing us to investigate different scenarios, with corresponding uncertainty.

13.7 Specifications for the regional changepoint model

In the regional changepoint model, each county has its own reproduction numbers, assumed constant in different periods, just like the national changepoint model. As there are more parameters in the regional changepoint model, we obtain 1000 parameter sets in the ABC-SMC.

Calibrating regional reproduction numbers is a more difficult estimation problem than calibrating national reproduction numbers, as we have a lot more parameters, and in addition less data in each county. Therefore, we cannot include the same amount of changepoints in the regional model as we can for the national model.

After we have obtained the estimated parameters, we run the model with these 1000 parameter sets again, from the beginning until today, plus three weeks into the future (or for an additional year). In this way, we obtain different trajectories of the future, allowing us to investigate different scenarios, with corresponding uncertainty.

14 Parameters used today

Figures 30 and 31 indicate which assumptions we make in our model, related to hospitalisation. We obtained data from the Norwegian Pandemiregister. These estimates will be regularly updated, on the basis of new data.




Figure 30: Hospital assumptions and parameters used before 1 August




Figure 31: Hospital assumptions and parameters used after 1 August

Table 15: Estimated parameters

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	Period
R0s	2.223	2.807	3.09	3.103	3.378	4.015	Until March 14
R1s	0.364	0.467	0.504	0.502	0.536	0.641	From 15 March to 19 April
R2s	0.04	0.49	0.619	0.615	0.764	1.074	From 20 April to 10 May
R3s	0.085	0.539	0.725	0.703	0.883	1.309	From 11 May to 30 June
R4s	0.003	0.653	0.883	0.888	1.146	1.799	From 01 July to 31 July
R5s	0.653	0.938	1.038	1.044	1.141	1.457	From 01 August to 31 August
R6s	0.686	0.855	0.916	0.922	0.989	1.127	From 01 September to 30 September
R7s	1.079	1.207	1.276	1.279	1.343	1.534	From 01 October to 25 October
R8s	1.05	1.201	1.277	1.279	1.348	1.57	From 26 October to 04 November
R9s	0.707	0.775	0.794	0.793	0.809	0.869	From 05 November to 30 November
R10s	1.002	1.06	1.079	1.079	1.098	1.156	From 01 December to 03 January
R11s	0.499	0.57	0.6	0.603	0.637	0.741	From 04 January to 21 January
R12s	0.615	0.812	0.852	0.849	0.895	1.099	From 22 January to 07 February
R13s	1.069	1.283	1.361	1.361	1.446	1.702	From 08 February to 21 February
R14s	1.164	1.336	1.407	1.404	1.462	1.666	From 22 February
AMPs	1.074	1.743	2.083	2.125	2.506	3.493	-
π_0	-0.795	-0.042	0.13	0.115	0.292	0.908	-
π_1	9.4e-08	1.2e-05	2.2e-05	2.7e-05	3.7e-05	8.9e-05	-
delays	1	3	3	3.115	4	4	-




Figure 32: Estimated densities of the reproduction numbers. National model

Table 16

R	Parameter	County	From	To	Pr(R>1)
5.66 (5.03-6.14)	R0	Oslo	2020-02-17	2020-03-14	1
3.5 (2.81-4.15)	R0	Rogaland	2020-02-17	2020-03-14	1
3.15 (1.28-4.54)	R0	Møre og Romsdal	2020-02-17	2020-03-14	1
3.3 (1.38-5.88)	R0	Nordland	2020-02-17	2020-03-14	0.98
3.61 (2.65-4.2)	R0	Viken	2020-02-17	2020-03-14	1
2.92 (1.69-4.82)	R0	Innlandet	2020-02-17	2020-03-14	1
3.43 (2.26-4.49)	R0	Vestfold og Telemark	2020-02-17	2020-03-14	1
2.77 (1.76-3.7)	R0	Agder	2020-02-17	2020-03-14	1
2.84 (1.91-4.12)	R0	Vestland	2020-02-17	2020-03-14	1
3.92 (2.4-5.16)	R0	Trøndelag	2020-02-17	2020-03-14	0.99
2.62 (1.43-3.92)	R0	Troms og Finnmark	2020-02-17	2020-03-14	1
0.59 (0.5-0.7)	R1	Oslo	2020-03-15	2020-04-19	0
0.48 (0.16-0.93)	R2	Oslo	2020-04-20	2020-07-24	0.02
1.31 (1.22-1.4)	R3	Oslo	2020-07-25	2020-09-30	1
1.5 (1.42-1.56)	R4	Oslo	2020-10-01	2020-11-04	1
1.03 (0.96-1.09)	R5	Oslo	2020-11-05	2020-12-14	0.8
1.38 (1.18-1.55)	R6	Oslo	2020-12-15	2021-01-03	1
0.78 (0.65-0.9)	R7	Oslo	2021-01-04	2021-02-04	0
1.77 (1.47-2.09)	R8	Oslo	2021-02-05	2021-02-21	1
1.31 (1.02-1.63)	R9	Oslo	2021-02-22		0.98
0.05 (0.01-0.12)	R1	Rogaland	2020-03-15	2020-04-19	0
0.78 (0.46-1.11)	R2	Rogaland	2020-04-20	2020-08-31	0.12
0.84 (0.62-1.02)	R3	Rogaland	2020-09-01	2020-11-04	0.04
0.62 (0.13-1.02)	R4	Rogaland	2020-11-05	2020-11-30	0.03
1.42 (1.17-1.71)	R5	Rogaland	2020-12-01	2021-01-03	1
0.17 (0.02-0.36)	R6	Rogaland	2021-01-04	2021-01-31	0
1.19 (0.6-1.6)	R7	Rogaland	2021-02-01		0.81
0.38 (0.04-0.66)	R1	Møre og Romsdal	2020-03-15	2020-04-19	0
0.64 (0.33-0.93)	R2	Møre og Romsdal	2020-04-20	2020-09-14	0.01
0.72 (0.34-1.09)	R3	Møre og Romsdal	2020-09-15	2020-11-04	0.07
0.5 (0.1-1.09)	R4	Møre og Romsdal	2020-11-05	2020-12-14	0.04
0.7 (0.1-1.51)	R5	Møre og Romsdal	2020-12-15	2021-01-03	0.24
0.4 (0.06-0.76)	R6	Møre og Romsdal	2021-01-04	2021-02-04	0
0.52 (0.04-1.16)	R7	Møre og Romsdal	2021-02-05		0.07
0.56 (0.11-0.87)	R1	Nordland	2020-03-15	2020-04-19	0
0.8 (0.51-1.07)	R2	Nordland	2020-04-20	2020-07-24	0.06
0.48 (0.02-1.43)	R3	Nordland	2020-07-25	2020-08-14	0.08
0.64 (0.3-1.04)	R4	Nordland	2020-08-15	2020-10-04	0.04
0.64 (0.27-1.09)	R5	Nordland	2020-10-05	2020-11-04	0.05
0.59 (0.03-1.19)	R6	Nordland	2020-11-05	2020-12-14	0.21
0.68 (0.01-1.51)	R7	Nordland	2020-12-15	2021-01-03	0.2
0.67 (0.22-1.09)	R8	Nordland	2021-01-04	2021-02-04	0.07
1.22 (0.75-1.65)	R9	Nordland	2021-02-05		0.84
0.29 (0.14-0.41)	R1	Viken	2020-03-15	2020-04-19	0
0.87 (0.42-1.14)	R2	Viken	2020-04-20	2020-07-24	0.29
1.09 (1-1.17)	R3	Viken	2020-07-25	2020-10-09	0.97
1.46 (1.38-1.53)	R4	Viken	2020-10-10	2020-11-04	1
0.82 (0.73-0.89)	R5	Viken	2020-11-05	2020-11-30	0
0.94 (0.85-1.01)	R6	Viken	2020-12-01	2021-01-03	0.04
0.88 (0.74-0.99)	R7	Viken	2021-01-04	2021-02-04	0.01
1.26 (1.04-1.49)	R8	Viken	2021-02-05	2021-02-21	0.99
1.3 (1.05-1.59)	R9	Viken	2021-02-22		0.99

Mean and 95% credible intervals

Table 17

R	Parameter	County	From	To	Pr(R>1)
0.5 (0-0.74)	R1	Innlandet	2020-03-15	2020-04-19	0
0.97 (0.74-1.18)	R2	Innlandet	2020-04-20	2020-07-24	0.49
0.84 (0.56-1.06)	R3	Innlandet	2020-07-25	2020-10-09	0.07
1.64 (1.12-2.05)	R4	Innlandet	2020-10-10	2020-10-24	0.99
0.5 (0.18-0.89)	R5	Innlandet	2020-10-25	2020-11-04	0
0.81 (0.34-1.14)	R6	Innlandet	2020-11-05	2020-12-14	0.17
0.48 (0.08-0.96)	R7	Innlandet	2020-12-15	2021-01-03	0.02
0.4 (0.07-0.73)	R8	Innlandet	2021-01-04	2021-02-04	0
0.48 (0.08-0.89)	R9	Innlandet	2021-02-05		0
0.19 (0.07-0.31)	R1	Vestfold og Telemark	2020-03-15	2020-04-19	0
0.95 (0.58-1.2)	R2	Vestfold og Telemark	2020-04-20	2020-07-24	0.32
0.76 (0.5-1.02)	R3	Vestfold og Telemark	2020-07-25	2020-10-09	0.03
0.81 (0.56-1.06)	R4	Vestfold og Telemark	2020-10-10	2020-11-04	0.07
1.05 (0.3-1.74)	R5	Vestfold og Telemark	2020-11-05	2020-11-19	0.55
0.93 (0.7-1.15)	R6	Vestfold og Telemark	2020-11-20	2021-01-03	0.21
0.58 (0.36-0.83)	R7	Vestfold og Telemark	2021-01-04	2021-02-04	0
1.46 (0.99-1.97)	R8	Vestfold og Telemark	2021-02-05	2021-02-21	0.97
1.68 (1.18-2.19)	R9	Vestfold og Telemark	2021-02-22		1
0.37 (0.1-0.57)	R1	Agder	2020-03-15	2020-04-19	0
0.9 (0.41-1.3)	R2	Agder	2020-04-20	2020-07-31	0.31
0.74 (0.36-1.07)	R3	Agder	2020-08-01	2020-09-19	0.04
0.88 (0.45-1.2)	R4	Agder	2020-09-20	2020-10-09	0.29
0.83 (0.23-1.4)	R5	Agder	2020-10-10	2020-11-04	0.24
0.47 (0.14-0.83)	R6	Agder	2020-11-05	2021-01-03	0
0.93 (0.75-1.1)	R7	Agder	2021-01-04	2021-02-11	0.24
1.13 (0.73-1.47)	R8	Agder	2021-02-12		0.76
0.42 (0.16-0.63)	R1	Vestland	2020-03-15	2020-04-19	0
0.8 (0.59-1.05)	R2	Vestland	2020-04-20	2020-07-24	0.06
1.36 (1.15-1.6)	R3	Vestland	2020-07-25	2020-09-04	1
0.96 (0.77-1.09)	R4	Vestland	2020-09-05	2020-10-09	0.28
1.68 (1.52-1.81)	R5	Vestland	2020-10-10	2020-11-04	1
0.41 (0.09-0.68)	R6	Vestland	2020-11-05	2020-11-30	0
0.57 (0.11-1.01)	R7	Vestland	2020-12-01	2021-01-03	0.03
0.96 (0.72-1.17)	R8	Vestland	2021-01-04	2021-02-04	0.37
0.74 (0.39-1.06)	R9	Vestland	2021-02-05		0.06
0.24 (0.04-0.5)	R1	Trøndelag	2020-03-15	2020-04-19	0
0.61 (0.26-0.97)	R2	Trøndelag	2020-04-20	2020-08-31	0.02
0.57 (0.21-1.05)	R3	Trøndelag	2020-09-01	2020-11-04	0.04
0.8 (0.19-1.46)	R4	Trøndelag	2020-11-05	2020-11-30	0.37
1.31 (0.94-1.54)	R5	Trøndelag	2020-12-01	2021-01-03	0.95
0.36 (0.13-0.57)	R6	Trøndelag	2021-01-04	2021-02-04	0
0.72 (0.1-1.3)	R7	Trøndelag	2021-02-05		0.22
0.17 (0.02-0.37)	R1	Troms og Finnmark	2020-03-15	2020-04-19	0
0.77 (0.07-1.23)	R2	Troms og Finnmark	2020-04-20	2020-09-14	0.28
0.72 (0.03-1.09)	R3	Troms og Finnmark	2020-09-15	2020-11-04	0.1
0.66 (0.07-1.31)	R4	Troms og Finnmark	2020-11-05	2020-11-30	0.18
0.54 (0.04-1.07)	R5	Troms og Finnmark	2020-12-01	2021-01-03	0.04
0.16 (0.01-0.44)	R6	Troms og Finnmark	2021-01-04	2021-02-04	0
1.25 (0.64-1.79)	R7	Troms og Finnmark	2021-02-05		0.82
1.25 (1.05-1.52)	AMP factor	All			-

Mean and 95% credible intervals

Table 18: Assumptions

Assumptions	Mean	Distribution	Reference
Mobile Mobility Data			
Telenor coverage	48%		https://ekomstatistikken.nkom.no/
Data updated	March 13th		
Data used in the predictions	March 12th	Fixed	Corrected to preserve population
Model parameters			
Exposed period ($1/\lambda_1$)	3 days	Exponential	Feretti et al 2020
Pre-symptomatic period ($1/\lambda_2$)	2 days	Exponential	Feretti et al 2020
Symptomatic infectious period ($1/\gamma$)	5 days	Exponential	Feretti et al 2020
Asymptomatic, infectious period ($1/\gamma$)	5 days	Exponential	Feretti et al 2020
Infectiousness asympt. (r_{I_a})	0.1	Fixed	Feretti et al 2020
Infectiousness presymp (r_{E_2})	1.25	Fixed	guided by Feretti et al 2020
Prob. asymptomatic infection (p_a)	0.4		Feretti et al 2020
Healthcare			
Time sympt. onset to hospitalisation	8.87 days (before August 1st) / 7.40 (After August 1st)	Neg. binomial	
Fraction asymptomatic infections	40%	Fixed	Mizumoto et al 2020 20% for the old population, Diamond Princess
% symptomatic and asymptomatic infections requiring hospitalization:			Salje et al 2020 corrected for: % of elderly living in elderly homes in Norway (last two age groups) and corrected for presence among positive tested since May 1.
0-9 years	0.1%	Fixed	
10 - 19 years	0.1%		
20 - 29 years	0.5%		
30 - 39 years	1.1%		
40 - 49 years	1.4%		
50 - 59 years	2.9%		
60 - 69 years	5.8%		
70 - 79 years	9.3%		
80+ years	22.3%		
% hospitalized patients requiring ICU			
Feb - July	15.1%	Fixed	Estimated from "Beredskapsregistret BereditC19"
August -	8.5%		
Probability that an admission has been reported on Monday			
From Sunday	32%	Fixed	Estimated from "Beredskapsregistret BereditC19"
From Saturday	49%		
From Friday	68%		
From Thursday	86%		
Probability that an admission has been reported			
From one day before	53%	Fixed	Estimated from "Beredskapsregistret BereditC19"
From two days before	77%		
From three days before	82%		
From four days before	91%		
Probability that a positive laboratory test has been reported			
From one day before	6.7%	Fixed	Estimated from MSIS
From two days before	59%		
From three days before	90%		
From four days before	97%		
Probability that a negative laboratory test has been reported			
From one day before	16%	Fixed	Estimated from MSIS
From two days before	74%		
From three days before	92%		
From four days before	98%		

Supplementary analysis: EpiEstim estimation of reproduction number based on laboratory-confirmed cases

To complement the results of the metapopulation model, we present estimates of the temporal evolution of the reproduction number in Norway based on an analysis of laboratory-confirmed cases. The primary purpose of this analysis is to provide a more comprehensive perspective on the epidemic situation, taking into account several data sources.

The hospitalisation data are a less biased information source for the number of infections compared to case data because the testing criteria in Norway has changed. For this reason, the present results should be interpreted with caution. During the early part of the period, testing of individuals was mainly based on travel history to areas with an ongoing outbreak. Since the middle of March, testing is recommended for people with an acute respiratory infection. From early May, the testing criteria have been expanded to include less severe symptoms. The analysis of laboratory-confirmed cases does not take into account the effect of imported cases during the early outbreak in Norway; the early results are less reliable than later results when the impact of importations is negligible.

EpiEstim method and assumptions: We estimate the instantaneous reproduction number using the procedure outlined in Thompson et al. (2019). This method, implemented in the EpiEstim R-package, uses a Bayesian approach to estimate the instantaneous reproduction number smoothed over a sliding window of 5 days, see figure 33. For the results to be comparable to those of the metapopulation model, we use the same natural history parameters. We estimate the date of infection for each confirmed case by first estimating the date of symptom onset and then subtracting 5 days for the incubation period. We estimate the date of symptom onset from the empirical delay between onset and testing in the first reported cases. For each case, we draw 100 possible onset dates from the delay distribution; this gives us 100 epi-curves that we use to estimate the reproduction number. The displayed results are the combined results from all these 100 simulated epi-curves. The serial interval was assumed to be 5 days with uncertainty; the serial interval refers to the time between symptom onset between successive cases in a chain of transmission (see <https://www.medrxiv.org/content/10.1101/2020.02.03.20019497v2>). To account for censoring of observations with onset dates in the last few days we correct the observed data by the mean of a negative binomial distribution with observation probability given by the empirical cumulative distribution of the onset to reporting date distributions. Due to this correction, the results from the last few days are uncertain, as indicated by increasing credible intervals.




Figure 33: Reproduction number estimated using the R package EpiEstim.

FHI COVID-19 modelling team:

- **Birgitte Freiesleben de Blasio** - Department of Method Development and Analytics. Norwegian Institute of Public Health and Oslo Centre for Biostatistics and Epidemiology, University of Oslo.
- **Francesco Di Ruscio** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Gunnar Øyvind Isaksson Rø** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Solveig Engebretsen** - Norsk Regnesentral.
- **Arnoldo Frigessi** - Oslo Centre for Biostatistics and Epidemiology, University of Oslo and Oslo University Hospital.
- **Alfonso Diz-Lois Palomares** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **David Swanson** - Oslo Centre for Biostatistics and Epidemiology, Oslo University Hospital.
- **Magnus Nygård Osnes** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Anja Bråthen Kristoffersen** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Kenth Engø-Monsen** - Telenor Research.
- **Louis Yat Hin Chan** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Jonas Christoffer Lindstrøm** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Richard White** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Gry Marysol Grøneng** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Chi Zhang** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Jørgen Eriksson Midtbø** - Department of Method Development and Analytics. Norwegian Institute of Public Health.